

Gust J, Wright JJ, Pratt EB and Bosma MM. Development of synchronized activity of cranial motor neurons in the segmented embryonic mouse hindbrain. *J. Physiol.* 550:123-133. 2003.

** Cover article

Corlew R, Bosma MM, Moody WJ. Spontaneous, synchronous electrical activity in neonatal mouse cortical neurones. *J. Physiol.* 560: 377-390. 2004.

Hunt PN, McCabe AK and Bosma MM. Midline serotonergic neurons drive widespread synchronized activity in embryonic mouse hindbrain. *J. Physiol.* 566:807-819. 2005.

** Cover article

** Perspectives article by Michael O'Donovan. Serotonergic neurones drive spontaneous activity in the developing mouse hindbrain. *J. Physiol.* 566:643. 2005.

Hunt PN, McCabe AK, Gust J and Bosma MM. Spatial restriction of spontaneous activity towards the rostral primary initiating zone during development of the embryonic mouse hindbrain. *J Neurobiol.* (now *Dev. Neurobiol.*) 66:1225-38. 2006.

Hunt PN, Gust J, McCabe AK, and Bosma MM. Primary role of the serotonergic midline system in synchronized spontaneous activity during development of the embryonic mouse hindbrain. *J Neurobiol.* (now *Dev. Neurobiol.*) 66:1239-52. 2006.

** Cover article

Hughes SM, Easton CR, Bosma MM. Properties and Mechanisms of Spontaneous Activity in the Embryonic Chick Hindbrain. *Dev. Neurobiol.* 69:477-90. 2009.

Rockhill W, Kirkman JL, Bosma MM. Spontaneous Activity in the Developing Mouse Midbrain Driven by an External Pacemaker. *Dev. Neurobiol.* 69: 689-704. 2009.

** Cover article

Moruzzi AM, Abedini NC, Hansen MA, Olson JE, Bosma MM. Differential expression of membrane conductances underlies spontaneous event initiation by rostral midline neurons in the embryonic mouse hindbrain. *J. Physiol.* 587:5081-5093. 2009.

** Perspectives article by Laura Ballerini. Insights into medio-lateral signaling in the developing mouse hindbrain: properties of midline drivers of network activity. *J. Physiol.* 587:5007. 2009.

Han S, Yu FH, Schwartz MD, Linton JD, Bosma MM, Hurley JB, Catterall WA, de la Iglesia HO. 2012. Na(V)1.1 channels are critical for intercellular communication in the suprachiasmatic nucleus and for normal circadian rhythms. *Proc Natl Acad Sci U S A.* 109:E368-77.

Watari H, Tose AJ, Bosma MM. 2013. Hyperpolarization of resting membrane potential causes retraction of spontaneous Ca^{2+} transients during mouse embryonic circuit development. *J Physiol.* 591:973-83.

Watari H, Tose AJ, Bosma MM. 2014. Looping circuit: a novel mechanism for prolonged spontaneous $[Ca^{2+}]_i$ increases in developing embryonic mouse brainstem. *J Physiol.* 592:711-27.